arXiv:1301.2705v1 [math.AG] 12 Jan 2013

HYPERGEOMETRIC SOLUTIONS OF THE QUANTUM DIFFERENTIAL EQUATION OF THE COTANGENT BUNDLE OF A PARTIAL FLAG VARIETY

V. TARASOV° AND A. VARCHENKO⋄

°*Department of Mathematical Sciences, Indiana University – Purdue University Indianapolis*
402 North Blackford St, Indianapolis, IN 46202-3216, USA

°*St. Petersburg Branch of Steklov Mathematical Institute*
Fontanka 27, St. Petersburg, 191023, Russia

⋄*Department of Mathematics, University of North Carolina at Chapel Hill*
Chapel Hill, NC 27599-3250, USA

Abstract. We describe hypergeometric solutions of the quantum differential equation of the cotangent bundle of a $\mathfrak{gl}_n$ partial flag variety. These hypergeometric solutions manifest the Landau-Ginzburg mirror symmetry for the cotangent bundle of a partial flag variety.

Contents

1. Introduction 2
2. Cotangent bundles of partial flag varieties 3
2.1. Partial flag varieties 3
2.2. Equivariant cohomology 4
3. Yangian 4
3.1. Yangian $Y(\mathfrak{gl}_N)$ 4
3.2. Algebra $\widetilde{Y}(\mathfrak{gl}_N)$ 5
3.3. $\widetilde{Y}(\mathfrak{gl}_N)$-action on $(\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}[\boldsymbol{z}; h]$ 5
3.4. $\widetilde{Y}(\mathfrak{gl}_N)$-action on $H^*_T(\mathcal{X}_n)$ according to [RTV] 6
4. Dynamical Hamiltonians and quantum multiplication 6
4.1. Dynamical Hamiltonians 6
4.2. Dynamical Hamiltonians $X^q_{\boldsymbol{\lambda},i}$ on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ 7
4.3. Quantum multiplication by divisors on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ 8
4.4. Dynamical Hamiltonians on $(\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}[\boldsymbol{z}; h]$ 8
4.5. qKZ difference connection 9
5. Yangian $\widetilde{Y}(\mathfrak{gl}_N)$ weight functions 10
5.1. Weight functions W_I 10
5.2. Weight functions $W_{\sigma,I}$ 10
5.3. Stable envelope map 10
5.4. The inverse map 11
5.5. The difference connection on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ 11
6. Four more connections 12

°*E-mail*: vt@math.iupui.edu, vt@pdmi.ras.ru, supported in part by NSF grant DMS-0901616
⋄*E-mail*: anv@email.unc.edu, supported in part by NSF grant DMS-1101508

6.1. Trigonometric dynamical connection on $\pi_{V_{\boldsymbol{\lambda}}}$ 12
6.2. Rational qKZ difference connection on $\pi_{V_{\boldsymbol{\lambda}}}$ 12
6.3. Relations between flat sections 13
6.4. Trigonometric KZ connection on $W_{\boldsymbol{\lambda}}$ 13
6.5. Rational dynamical difference connection on $\pi_{W_{\boldsymbol{\lambda}}}$ 14
6.6. Equivalence of connections on $\pi_{V_{\boldsymbol{\lambda}}}$ and $\pi_{W_{\boldsymbol{\lambda}}}$ 14
7. Integral representations for flat sections 15
7.1. Master function 15
7.2. $\mathfrak{gl}_n$ weight functions 15
7.3. Hypergeometric integrals 16
7.4. Example 17
References 18

1. Introduction

In [MO], D. Maulik and A. Okounkov develop a general theory connecting quantum groups and equivariant quantum cohomology of Nakajima quiver varieties, see [N1, N2]. In particular, in [MO] the operators of quantum multiplication by divisors are described. As well-known, these operators determine the equivariant quantum differential equation of a quiver variety. In this paper we apply this description to the cotangent bundles of $\mathfrak{gl}_n$ partial flag varieties and construct hypergeometric solutions of the associated equivariant quantum differential equation.

In [GRTV] and [RTV], the equivariant quantum differential equation of the cotangent bundle of a $\mathfrak{gl}_n$ partial flag variety was identified with the trigonometric dynamical differential equation introduced in [TV4]. By the $(\mathfrak{gl}_N, \mathfrak{gl}_n)$-duality of [TV4], the trigonometric dynamical differential equation is identified with the trigonometric KZ differential equation. Hypergeometric solutions of the trigonometric KZ differential equation were constructed in [SV, MV]. By using this sequence of isomorphisms we obtain hypergeometric solutions of the equivariant quantum differential equation. The hypergeometric solutions have the form

$$I_\psi(\boldsymbol{z};\tilde{\boldsymbol{q}};h;\kappa) = \int_{\psi(\boldsymbol{z};\tilde{\boldsymbol{q}};h;\kappa)} \Phi(\boldsymbol{s};\boldsymbol{z};\tilde{\boldsymbol{q}};h)^{h/\kappa}\omega(\boldsymbol{s};\boldsymbol{z};\tilde{\boldsymbol{q}};h)d\boldsymbol{s}, \tag{1.1}$$

where $\boldsymbol{z} = (z_1, \dots, z_n)$, h are equivariant parameters, $\tilde{\boldsymbol{q}} = (\tilde{q}_1, \dots, \tilde{q}_N)$ quantum parameters, $\boldsymbol{s} = (s_j^{(i)})$ integration variables, κ the parameter of the differential equation, $\psi(\boldsymbol{z};\tilde{\boldsymbol{q}};h;\kappa)$ the integration cycle in the $\boldsymbol{s}$-space, $\omega(\boldsymbol{s};\boldsymbol{z};\tilde{\boldsymbol{q}};h)$ a rational function, $\Phi(\boldsymbol{s};\boldsymbol{z};\tilde{\boldsymbol{q}};h)$ the master function, see Corollary 7.4.

Studying solutions of the quantum differential equation may lead to better understanding Gromow-Witten invariants of the cotangent bundle of a partial flag variety, c.f. Givental's study of the J-function in [G1, G2, G3].

The existence of solutions of the quantum differential equation as such oscillatory integrals manifests the Landau-Ginzburg mirror symmetry for the cotangent bundle of a partial flag variety. One may think that the logarithm of the master function is the Landau-Ginzburg potential of the mirror dual object. In particular, one may expect that the algebra of functions on the critical set of the master function $\Phi(\,\cdot\,;\boldsymbol{z};\tilde{\boldsymbol{q}};h)$ is isomorphic to the corresponding localization of the equivariant quantum cohomology algebra of the cotangent bundle of a partial flag variety, see Section 7.4 and similar statements for Bethe algebras in [MTV2, MTV3].

The trigonometric KZ differential equation and trigonometric dynamical differential equation come in pairs with compatible difference equations. The trigonometric KZ differential equation is compatible with the rational dynamical difference equation introduced in [TV3]. The trigonometric dynamical differential equation is compatible with the rational *qKZ* difference equation, as shown in [TV4]. Under the $(\mathfrak{gl}_N, \mathfrak{gl}_n)$-duality, the pair consisting of the trigonometric KZ differential equation and rational dynamical difference equation is identified with the pair consisting of the trigonometric dynamical differential equation and rational *qKZ* difference equation, see [TV4]. Under the identification of the trigonometric dynamical differential equation with the equivariant quantum differential equation of [BMO], the rational *qKZ* difference operators are identified with the shift operators of [MO], see also [BMO].

It was shown in [MV], that the hypergeometric solutions for the trigonometric KZ differential equation also satisfy the difference dynamical equation. By using all of the above identifications, we conclude that the hypergeometric solutions (1.1) of the equivariant quantum differential equation give also flat sections of the difference connection defined by the shift operators of [MO] on the equivariant quantum cohomology of the cotangent bundle of a partial $\mathfrak{gl}_n$ flag variety.

As shown in [TV1, TV2], the *qKZ* equation has solutions in the form of multidimensional q-hypergeometric integrals. These q-hypergeometric integrals are expected to satisfy the compatible dynamical differential equation. That will imply that the quantum differential equation of [MO] has solutions in the form of multidimensional q-hypergeometric integrals. See an example of such solutions in Section 8.4 of [GRTV]. We plan to develop these q-hypergeometric solutions in a separate paper.

2. Cotangent bundles of partial flag varieties

2.1. **Partial flag varieties.** Fix natural numbers N, n. Let $\boldsymbol{\lambda} \in \mathbb{Z}^N_{\geqslant 0},\ |\boldsymbol{\lambda}| = \lambda_1 + \dots + \lambda_N = n$. Consider the partial flag variety $\mathcal{F}_{\boldsymbol{\lambda}}$ parametrizing chains of subspaces

$$0 \,=\, F_0 \subset F_1 \subset \ldots \subset F_N \,=\, \mathbb{C}^n$$

with $\dim F_i/F_{i-1} = \lambda_i,\ i = 1, \ldots, N$. Denote by $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ the cotangent bundle of $\mathcal{F}_{\boldsymbol{\lambda}}$. Denote

$$\mathcal{X}_n = \cup_{|\boldsymbol{\lambda}|=n} T^*\mathcal{F}_{\boldsymbol{\lambda}}.$$

Example. If $n = 1$, then $\boldsymbol{\lambda} = (0, \ldots, 0, 1_i, 0, \ldots, 0)$, $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ is a point and $\mathcal{X}_1$ is the union of N points.

If $n = 2$ then $\boldsymbol{\lambda} = (0, \ldots, 0, 1_i, 0, \ldots, 0, 1_j, 0, \ldots, 0)$ or $\boldsymbol{\lambda} = (0, \ldots, 0, 2_i, 0, \ldots, 0)$. In the first case $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ is the cotangent bundle of projective line, in the second case $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ is a point. Thus $\mathcal{X}_2$ is the union of N points and $N(N-1)/2$ copies of the cotangent bundle of projective line.

Let $I = (I_1, \ldots, I_N)$ be a partition of $\{1, \ldots, n\}$ into disjoint subsets $I_1, \ldots, I_N$. Denote $\mathcal{I}_{\boldsymbol{\lambda}}$ the set of all partitions I with $|I_j| = \lambda_j,\ j = 1, \ldots N$. Denote $\mathcal{I}_n = \cup_{|\boldsymbol{\lambda}|=n} \mathcal{I}_{\boldsymbol{\lambda}}$.

Let $u_1, \ldots, u_n$ be the standard basis of $\mathbb{C}^n$. For any $I \in \mathcal{I}_{\boldsymbol{\lambda}}$, let $x_I \in \mathcal{F}_{\boldsymbol{\lambda}}$ be the point corresponding to the coordinate flag $F_1 \subset \cdots \subset F_N$, where F_i is the span of the standard basis vectors $u_j \in \mathbb{C}^n$ with $j \in I_1 \cup \ldots \cup I_i$. We embed $\mathcal{F}_{\boldsymbol{\lambda}}$ in $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ as the zero section and consider the points x_I as points of $T^*\mathcal{F}_{\boldsymbol{\lambda}}$.

2.2. **Equivariant cohomology.** Denote $G = GL_n(\mathbb{C}) \times \mathbb{C}^\times$. Let $A \subset GL_n(\mathbb{C})$ be the torus of diagonal matrices. Denote $T = A \times \mathbb{C}^\times$ the subgroup of G.

The groups $A \subset GL_n$ act on $\mathbb{C}^n$ and hence on $T^*\mathcal{F}_{\boldsymbol{\lambda}}$. Let the group $\mathbb{C}^\times$ act on $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ by multiplication in each fiber. We denote by $-h$ its $\mathbb{C}^\times$-weight.

We consider the equivariant cohomology algebras $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}};\mathbb{C})$ and

$$H^*_T(\mathcal{X}_n) = \oplus_{|\boldsymbol{\lambda}|=n} H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}};\mathbb{C}).$$

Denote by $\Gamma_i = \{\gamma_{i,1},\ldots,\gamma_{i,\lambda_i}\}$ the set of the Chern roots of the bundle over $\mathcal{F}_{\boldsymbol{\lambda}}$ with fiber F_i/F_{i-1}. Let $\boldsymbol{\Gamma} = (\Gamma_1;\ldots;\Gamma_N)$. Denote by $\boldsymbol{z} = \{z_1,\ldots,z_n\}$ the Chern roots corresponding to the factors of the torus T. Then

$$H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}}) = \mathbb{C}[\boldsymbol{\Gamma}]^{S_{\lambda_1}\times\cdots\times S_{\lambda_N}} \otimes \mathbb{C}[\boldsymbol{z}] \otimes \mathbb{C}[h] \Big/ \Big\langle \prod_{i=1}^{N}\prod_{j=1}^{\lambda_i}(u-\gamma_{i,j}) = \prod_{a=1}^{n}(u-z_a)\Big\rangle .$$

The cohomology $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ is a module over $H^*_T(pt;\mathbb{C}) = \mathbb{C}[\boldsymbol{z}]\otimes\mathbb{C}[h]$.

Example. If $n=1$, then

$$H^*_T(\mathcal{X}_1) = \oplus_{i=1}^N H^*_T(T^*\mathcal{F}_{(0,\ldots,0,1_i,0,\ldots,0)})$$

is naturally isomorphic to $\mathbb{C}^N\otimes\mathbb{C}[z_1;h]$ with basis $v_i = (0,\ldots,0,1_i,0,\ldots,0)$, $i=1,\ldots,N$.

For $i=1,\ldots,N$, denote $\lambda^{(i)} = \lambda_1+\cdots+\lambda_i$. Denote $\Theta_i = \{\theta_{i,1},\ldots,\theta_{i,\lambda^{(i)}}\}$ the Chern roots of the bundle $\boldsymbol{F}_i$ over $\mathcal{F}_{\boldsymbol{\lambda}}$ with fiber F_i. Let $\boldsymbol{\Theta} = (\Theta_1,\ldots,\Theta_N)$. The relations

$$\prod_{j=1}^{\lambda^{(i)}}(u-\theta_{i,j}) = \prod_{\ell=1}^{i}\prod_{j=1}^{\lambda_i}(u-\gamma_{i,j}), \qquad i=1,\ldots,N,$$

define the homomorphism

$$\mathbb{C}[\boldsymbol{\Theta}]^{S_{\lambda^{(1)}}\times\cdots\times S_{\lambda^{(N)}}}\otimes\mathbb{C}[\boldsymbol{z}]\otimes\mathbb{C}[h] \to H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}}).$$

3. Yangian

3.1. **Yangian $Y(\mathfrak{gl}_N)$.** The Yangian $Y(\mathfrak{gl}_N)$ is the unital associative algebra with generators $T^{\{s\}}_{i,j}$ for $i,j=1,\ldots,N$, $s\in\mathbb{Z}_{>0}$, subject to relations

$$(u-v)\big[T_{i,j}(u),T_{k,l}(v)\big] = T_{k,j}(v)T_{i,l}(u) - T_{k,j}(u)T_{i,l}(v), \qquad i,j,k,l=1,\ldots,N,$$

where $T_{i,j}(u) = \delta_{i,j} + \sum_{s=1}^\infty T^{\{s\}}_{i,j}u^{-s}$. The Yangian $Y(\mathfrak{gl}_N)$ is a Hopf algebra with the coproduct $\Delta : Y(\mathfrak{gl}_N)\to Y(\mathfrak{gl}_N)\otimes Y(\mathfrak{gl}_N)$ given by $\Delta\big(T_{i,j}(u)\big) = \sum_{k=1}^N T_{k,j}(u)\otimes T_{i,k}(u)$ for $i,j=1,\ldots,N$. The Yangian $Y(\mathfrak{gl}_N)$ contains, as a Hopf subalgebra, the universal enveloping algebra $U(\mathfrak{gl}_N)$ of the Lie algebra $\mathfrak{gl}_N$. The embedding is given by $e_{i,j}\mapsto T^{\{1\}}_{j,i}$, where $e_{i,j}$ are standard standard generators of $\mathfrak{gl}_N$.

Notice that $\big[T^{\{1\}}_{i,j},T^{\{s\}}_{k,l}\big] = \delta_{i,l}T^{\{s\}}_{k,j} - \delta_{j,k}T^{\{s\}}_{i,l}$ for $i,j,k,l=1,\ldots,N$, $s\in\mathbb{Z}_{>0}$, which implies that the Yangian $Y(\mathfrak{gl}_N)$ is generated by the elements $T^{\{1\}}_{i,i+1}, T^{\{1\}}_{i+1,i}$, $i=1,\ldots,N-1$, and $T^{\{s\}}_{1,1}$, $s>0$.

3.2. **Algebra $\widetilde{Y}(\mathfrak{gl}_N)$.** In this section we follow [GRTV, Section 3.3]. In formulas of that Section 3.3 we replace h with $-h$.

Let $\widetilde{Y}(\mathfrak{gl}_N)$ be the subalgebra of $Y(\mathfrak{gl}_N)\otimes\mathbb{C}[h]$ generated over $\mathbb{C}$ by $\mathbb{C}[h]$ and the elements $(-h)^{s-1}T_{i,j}^{\{s\}}$ for $i,j=1,\dots,N,\ \ s>0$. Equivalently, the subalgebra $\widetilde{Y}(\mathfrak{gl}_N)$ is generated over $\mathbb{C}$ by $\mathbb{C}[h]$ and the elements $T_{i,i+1}^{\{1\}},\,T_{i+1,i}^{\{1\}},\ \ i=1,\dots,N-1$, and $(-h)^{s-1}T_{1,1}^{\{s\}},\ \ s>0$.

For $p=1,\dots,N,\ \ \boldsymbol{i}=\{1\leqslant i_1<\dots<i_p\leqslant N\},\ \ \boldsymbol{j}=\{1\leqslant j_1<\dots<j_p\leqslant N\}$, define

$$M_{\boldsymbol{i},\boldsymbol{j}}(u)=\sum_{\sigma\in S_p}(-1)^\sigma\,T_{i_1,j_{\sigma(1)}}(u)\dots T_{i_p,j_{\sigma(p)}}(u-p+1)\,.$$

Introduce the series $A_1(u),\dots,A_N(u),\ E_1(u),\dots,E_{N-1}(u),\ F_1(u),\dots,F_{N-1}(u)$:

$$A_p(u)\,=\,M_{\boldsymbol{i},\boldsymbol{i}}(-u/h)\,=\,1+\sum_{s=1}^\infty(-h)^s A_{p,s}\,u^{-s}\,,\tag{3.1}$$

$$E_p(u)\,=\,-h^{-1}M_{\boldsymbol{j},\boldsymbol{i}}(-u/h)\big(M_{\boldsymbol{i},\boldsymbol{i}}(-u/h)\big)^{-1}\,=\,\sum_{s=1}^\infty(-h)^{s-1}E_{p,s}\,u^{-s}\,,\tag{3.2}$$

$$F_p(u)\,=\,-h^{-1}\big(M_{\boldsymbol{i},\boldsymbol{i}}(-u/h)\big)^{-1}M_{\boldsymbol{i},\boldsymbol{j}}(-u/h)\,=\,\sum_{s=1}^\infty(-h)^{s-1}F_{p,s}\,u^{-s}\,,$$

where in formulas (3.1) and (3.2) we have $\boldsymbol{i}=\{1,\dots,p\}$, $\boldsymbol{j}=\{1,\dots,p-1,p+1\}$. Observe that $E_{p,1}=T_{p+1,p}^{\{1\}}$, $F_{p,1}=T_{p,p+1}^{\{1\}}$ and $A_{1,s}=T_{1,1}^{\{s\}}$, so the coefficients of the series $E_p(u)$, $F_p(u)$ and $h^{-1}\big(A_p(u)-1\big)$ together with $\mathbb{C}[h]$ generate $\widetilde{Y}(\mathfrak{gl}_N)$. In what follows we will describe actions of the algebra $\widetilde{Y}(\mathfrak{gl}_N)$ by using series (3.1), (3.2).

3.3. **$\widetilde{Y}(\mathfrak{gl}_N)$-action on $(\mathbb{C}^N)^{\otimes n}\otimes\mathbb{C}[\boldsymbol{z};h]$.** Set

$$L(u)\,=\,(u-z_n-hP^{(0,n)})\dots(u-z_1-hP^{(0,1)})\,,$$

where the factors of $\mathbb{C}^N\otimes(\mathbb{C}^N)^{\otimes n}$ are labeled by $0,1,\dots,n$ and $P^{(i,j)}$ is the permutation of the i-th and j-th factors. The operator $L(u)$ is a polynomial in $u,\boldsymbol{z},h$ with values in $\mathrm{End}(\mathbb{C}^N\otimes(\mathbb{C}^N)^{\otimes n})$. We consider $L(u)$ as an $N\times N$ matrix with $\mathrm{End}(V)\otimes\mathbb{C}[u;\boldsymbol{z};h]$-valued entries $L_{i,j}(u)$.

Proposition 3.1 (Proposition 4.1 in [GRTV]). *The assignment*

$$\phi\big(T_{i,j}(-u/h)\big)\,=\,L_{i,j}(u)\prod_{a=1}^n(u-z_a)^{-1}\tag{3.3}$$

defines the action of the algebra $\widetilde{Y}(\mathfrak{gl}_N)$ on $(\mathbb{C}^N)^{\otimes n}\otimes\mathbb{C}[\boldsymbol{z};h]$. Here the right-hand side of (3.3) *is a series in u^{-1} with coefficients in* $\mathrm{End}((\mathbb{C}^N)^{\otimes n})\otimes\mathbb{C}[\boldsymbol{z};h])$.

Under this action, the subalgebra $U(\mathfrak{gl}_N)\subset\widetilde{Y}(\mathfrak{gl}_N)$ acts on $(\mathbb{C}^N)^{\otimes n}\otimes\mathbb{C}[\boldsymbol{z};h]$ in the standard way: any element $x\in\mathfrak{gl}_N$ acts as $x^{(1)}+\dots+x^{(n)}$. The action ϕ was denoted in [GRTV] by ϕ^+.

3.4. **$\widetilde{Y}(\mathfrak{gl}_N)$-action on $H^*_T(\mathcal{X}_n)$ according to** [RTV]**.** We define the $\widetilde{Y}(\mathfrak{gl}_N)$-action ρ on $H^*_T(\mathcal{X}_n)$ by formulas (3.4), (3.6), (3.7) below. We define $\rho\bigl(A_p(u)\bigr) : H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda}) \to H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda})$ by

$$\rho\bigl(A_p(u)\bigr) \,:\, [f] \,\mapsto\, \Bigl[\,f(\boldsymbol\Gamma;\boldsymbol z;h)\,\prod_{a=1}^{p}\,\prod_{i=1}^{\lambda_p}\Bigl(1-\frac{h}{u-\gamma_{p,i}}\Bigr)\Bigr]\,, \tag{3.4}$$

for $p=1,\dots,N$. In particular,

$$\rho(X^\infty_i) \,:\, [f] \,\mapsto\, [(\gamma_{i,1}+\ldots+\gamma_{i,\lambda_i})\,f(\boldsymbol\Gamma;\boldsymbol z;h)]\,, \qquad i=1,\dots,N\,. \tag{3.5}$$

Let $\alpha_1,\dots,\alpha_{N-1}$ be simple roots, $\alpha_p=(0,\dots,0,1,-1,0,\dots,0)$, with $p-1$ first zeros. We define

$$\rho\bigl(E_p(u)\bigr) \,:\, H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda-\alpha_p}) \,\mapsto\, H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda})\,,$$

$$\rho\bigl(E_p(u)\bigr) \,:\, [f] \,\mapsto\, \Bigl[\,\sum_{i=1}^{\lambda_p}\frac{f(\boldsymbol\Gamma'^i;\boldsymbol z;h)}{u-\gamma_{p,i}}\,\prod_{\substack{j=1\\ j\neq i}}^{\lambda_p}\frac{1}{\gamma_{p,j}-\gamma_{p,i}}\,\prod_{k=1}^{\lambda_{p+1}}(\gamma_{p,i}-\gamma_{p+1,k}-h)\,\Bigr]\,, \tag{3.6}$$

$$\rho\bigl(F_p(u)\bigr) \,:\, H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda+\alpha_p}) \,\mapsto\, H^*_T(T^*\mathcal{F}_{\boldsymbol\lambda})\,,$$

$$\rho\bigl(F_p(u)\bigr) \,:\, [f] \,\mapsto\, \Bigl[\,\sum_{i=1}^{\lambda_{p+1}}\frac{f(\boldsymbol\Gamma^{i\prime};\boldsymbol z;h)}{u-\gamma_{p+1,i}}\,\prod_{\substack{j=1\\ j\neq i}}^{\lambda_{p+1}}\frac{1}{\gamma_{p+1,i}-\gamma_{p+1,j}}\,\prod_{k=1}^{\lambda_p}(\gamma_{p,k}-\gamma_{p+1,i}-h)\,\Bigr]\,, \tag{3.7}$$

where

$$\boldsymbol\Gamma'^i \,=\, (\Gamma_1;\ldots;\Gamma_{p-1};\Gamma_p-\{\gamma_{p,i}\};\Gamma_{p+1}\cup\{\gamma_{p,i}\};\Gamma_{p+2};\ldots;\Gamma_N)\,,$$

$$\boldsymbol\Gamma^{i\prime} \,=\, (\Gamma_1;\ldots;\Gamma_{p-1};\Gamma_p\cup\{\gamma_{p+1,i}\};\Gamma_{p+1}-\{\gamma_{p+1,i}\};\Gamma_{p+2};\ldots;\Gamma_N)\,.$$

Theorem 3.2 (Theorem 5.10 in [GRTV])**.** *These formulas define a $\widetilde{Y}(\mathfrak{gl}_N)$-module structure on $H^*_T(\mathcal{X}_n)$.*

This $\widetilde{Y}(\mathfrak{gl}_N)$-module structure was denoted in [GRTV] by ρ^- and h in [GRTV] is replaced with $-h$. The topological interpretation of this $\widetilde{Y}(\mathfrak{gl}_N)$-action see in [GRTV, Theorem 5.16].

In [MO], a Yangian module structure on $H^*_T(\mathcal{X}_n)$ was introduced.

Theorem 3.3 (Corollary 6.4 in [RTV])**.** *The $\widetilde{Y}(\mathfrak{gl}_N)$-module structure ρ on $H^*_T(\mathcal{X}_n)$ coincides with the Yangian module structure on $H^*_T(\mathcal{X}_n)$ introduced in* [MO].

4. Dynamical Hamiltonians and quantum multiplication

4.1. **Dynamical Hamiltonians.** Assume that $q_1,\dots,q_N$ are distinct numbers. Define the elements $X^q_1,\dots,X^q_N \in \widetilde{Y}(\mathfrak{gl}_N)$ by the rule

$$X^q_i \,=\, -hT^{\{2\}}_{i,i}+\frac{h}{2}\,e_{i,i}\bigl(e_{i,i}-1\bigr)\,-\,h\sum_{\substack{j=1\\ j\neq i}}^{N}\frac{q_j}{q_i-q_j}\,G_{i,j}\,,$$

where $G_{i,j} = e_{i,j}e_{j,i} - e_{i,i} = e_{j,i}e_{i,j} - e_{j,j}$. By taking the limit $q_{i+1}/q_i \to 0$ for all $i = 1, \dots, N-1$, we define the elements $X_1^\infty, \dots, X_N^\infty \in \widetilde{Y}(\mathfrak{gl}_N)$,

$$X_i^\infty = -hT_{i,i}^{\{2\}} + \frac{h}{2}\, e_{i,i}\big(e_{i,i} - 1\big) + h\,(G_{i,1} + \ldots + G_{i,i-1})\,,$$

see [GRTV]. The elements X_i^q, X_i^∞, $i = 1, \dots, N$, are called the dynamical Hamiltonians. Observe that

$$X_i^q = X_i^\infty - h\sum_{j=1}^{i-1} \frac{q_i}{q_i - q_j}\, G_{i,j} - h\sum_{j=i+1}^{n} \frac{q_j}{q_i - q_j}\, G_{i,j}\,.$$

Given $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$, set $G_{\boldsymbol{\lambda},i,j} = e_{j,i}e_{i,j}$ for $\lambda_i \geqslant \lambda_j$ and $G_{\boldsymbol{\lambda},i,j} = e_{i,j}e_{j,i}$ for $\lambda_i < \lambda_j$. We define the elements $X_{\boldsymbol{\lambda},1}^q, \dots, X_{\boldsymbol{\lambda},N}^q \in \widetilde{Y}(\mathfrak{gl}_N)$,

$$X_{\boldsymbol{\lambda},i}^q = X_i^\infty - h\sum_{j=1}^{i-1} \frac{q_i}{q_i - q_j}\, G_{\boldsymbol{\lambda},i,j} - h\sum_{j=i+1}^{n} \frac{q_j}{q_i - q_j}\, G_{\boldsymbol{\lambda},i,j}\,.$$

Let $\kappa \in \mathbb{C}^\times$. The formal differential operators

$$\nabla_{\boldsymbol{q},\kappa,i} = \kappa\, q_i \frac{\partial}{\partial q_i} - X_i^q, \qquad i = 1, \dots, N, \tag{4.1}$$

pairwise commute and, hence, define a flat connection $\nabla_{\boldsymbol{q},\kappa}$ for any $\widetilde{Y}(\mathfrak{gl}_N)$-module, see [GRTV].

Lemma 4.1 (Lemma 3.5 in [GRTV]). *The connection* $\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa}$ *defined by*

$$\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa,i} = \kappa\, q_i \frac{\partial}{\partial q_i} - X_{\boldsymbol{\lambda},i}^q\,, \tag{4.2}$$

$i = 1, \dots, N$, *is flat for any* κ.

Proof. The connection $\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa}$ is gauge equivalent to connection ∇_κ,

$$\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa,i} = (\Upsilon_{\boldsymbol{\lambda}})^{-1}\, \nabla_{\boldsymbol{q},\kappa,i}\, \Upsilon_{\boldsymbol{\lambda}}\,, \qquad \Upsilon_{\boldsymbol{\lambda}} = \prod_{1 \leqslant i < j \leqslant N} (1 - q_j/q_i)^{h\varepsilon_{\boldsymbol{\lambda},i,j}/\kappa}\,, \tag{4.3}$$

where $\varepsilon_{\boldsymbol{\lambda},i,j} = e_{j,j}$ for $\lambda_i \geqslant \lambda_j$, and $\varepsilon_{\boldsymbol{\lambda},i,j} = e_{i,i}$ for $\lambda_i < \lambda_j$. □

Connection (4.1) was introduced in [TV4], see also Appendix B in [MTV1], and is called the *trigonometric dynamical connection*. Later the definition was extended from $\mathfrak{sl}_N$ to other simple Lie algebras in [TL] under the name of the trigonometric Casimir connection.

The trigonometric dynamical connection is defined over $\mathbb{C}^N$ with coordinates $q_1, \dots, q_N$, it has singularities at the union of the diagonals $q_i = q_j$.

4.2. **Dynamical Hamiltonians $X_{\boldsymbol{\lambda},i}^q$ on $H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$.** Recall the $\widetilde{Y}(\mathfrak{gl}_N)$-module structure ρ defined on $H_T^*(\mathcal{X}_n) = \oplus_{|\boldsymbol{\lambda}|=n} H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ in Section 3.4. For any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_N) \in \mathbb{Z}_{\geqslant 0}^N$, $|\boldsymbol{\mu}| = n$, the action of the dynamical Hamiltonians $X_{\boldsymbol{\mu},i}^q$ preserve each of $H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$.

Lemma 4.2 (Lemma 7.6 in [RTV])**.** *For any* $\boldsymbol{\lambda}$ *and* $i = 1, \dots, n$*, the restriction of* $\rho(X^q_{\boldsymbol{\lambda},i})$ *to* $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *has the form:*

(4.4)

$$\begin{aligned}\rho(X^q_{\boldsymbol{\lambda},i}) \;&=\; (\gamma_{i,1} + \dots + \gamma_{i,\lambda_i}) - h\sum_{j=1}^{i-1} \frac{q_i}{q_i - q_j}\,\rho(G_{\boldsymbol{\lambda},i,j}) - h\sum_{j=i+1}^{n} \frac{q_j}{q_i - q_j}\,\rho(G_{\boldsymbol{\lambda},i,j}) = \\ &=\; (\gamma_{i,1} + \dots + \gamma_{i,\lambda_i}) - h\sum_{j=1}^{i-1} \frac{q_i}{q_i - q_j}\,\rho(e_{j,i}e_{i,j}) - h\sum_{j=i+1}^{n} \frac{q_j}{q_i - q_j}\,\rho(e_{i,j}e_{j,i}) \,+ C,\end{aligned}$$

where $(\gamma_{i,1} + \dots + \gamma_{i,\lambda_i})$ *denotes the operator of multiplication by the cohomology class* $\gamma_{i,1} + \dots + \gamma_{i,\lambda_i}$*, the operator* C *is a scalar operator on* $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$*, and for any* $i \neq j$ *the element* $\rho(G_{\boldsymbol{\lambda},i,j})$ *annihilates the identity element* $1_{\boldsymbol{\lambda}} \in H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$*.*

4.3. **Quantum multiplication by divisors on** $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$**.** In [MO], the quantum multiplication by divisors on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ is described. The fundamental equivariant cohomology classes of divisors on $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ are linear combinations of $D_i = \gamma_{i,1} + \dots + \gamma_{i,\lambda_i}$, $i = 1, \dots, N$. The quantum multiplication $D_i *_{\tilde{\boldsymbol{q}}}$ depends on parameters $\tilde{\boldsymbol{q}} = (\tilde q_1, \dots, \tilde q_N) \in (\mathbb{C}^\times)^N$.

Theorem 4.3 (Theorem 10.2.1 in [MO])**.** *For* $i = 1, \dots, N$*, the quantum multiplication by* D_i *is given by the formula:*

(4.5)

$$D_i *_{\tilde{\boldsymbol{q}}} \;=\; (\gamma_{i,1} + \dots + \gamma_{i,\lambda_i}) + h\sum_{j=1}^{i-1} \frac{\tilde q_j/\tilde q_i}{1 - \tilde q_j/\tilde q_i}\,\rho(e_{j,i}e_{i,j}) - h\sum_{j=i+1}^{n} \frac{\tilde q_i/\tilde q_j}{1 - \tilde q_i/\tilde q_J}\,\rho(e_{i,j}e_{j,i}) \,+ C,$$

where C *is a scalar operator on* $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *fixed by the requirement that the purely quantum part of* $D_i *_{\tilde{\boldsymbol{q}}}$ *annihilates the identity* $1_{\boldsymbol{\lambda}}$*.*

Corollary 4.4 (Corollary 7.8 in [RTV])**.** *For* $i = 1, \dots, N$*, the operator* $D_i *_{\tilde{\boldsymbol{q}}}$ *of quantum multiplication by* D_i *on* $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *equals the action* $\rho(X^q_{\boldsymbol{\lambda},i})$ *on* $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *of the dynamical Hamiltonian* $X^q_{\boldsymbol{\lambda},i}$ *if we put* $(q_1, \dots, q_N) = (\tilde q_1^{-1}, \dots, \tilde q_N^{-1})$*.*

The quantum connection $\nabla_{\mathrm{quant},\boldsymbol{\lambda},\tilde{\boldsymbol{q}},\kappa}$ on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ is defined by the formula

$$\nabla_{\mathrm{quant},\boldsymbol{\lambda},\tilde{\boldsymbol{q}},\kappa,i} \;=\; \kappa\,\tilde q_i \frac{\partial}{\partial \tilde q_i} \,-\, D_i *_{\tilde{\boldsymbol{q}}}\,, \qquad i = 1, \dots, N, \tag{4.6}$$

where $\kappa \in \mathbb{C}^\times$ is a parameter of the connection, see [BMO]. By Corollary 4.4, we have

$$\nabla_{\mathrm{quant},\boldsymbol{\lambda},\tilde{\boldsymbol{q}},\kappa,i} = \rho(\nabla_{\boldsymbol{\lambda},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa}), \qquad i = 1, \dots, N. \tag{4.7}$$

4.4. **Dynamical Hamiltonians on** $(\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}[\boldsymbol{z}; h]$**.** Recall that $e_{i,j}$, $i, j = 1, \dots, N$, denote standard generators of $\mathfrak{gl}_N$. A vector v of a $\mathfrak{gl}_N$-module M has weight $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N) \in \mathbb{C}^N$ if $e_{i,i}v = \lambda_i v$ for $i = 1, \dots, N$. We denote by $M_{\boldsymbol{\lambda}} \subset M$ the weight subspace of weight $\boldsymbol{\lambda}$.

We consider $\mathbb{C}^N$ as the standard vector representation of $\mathfrak{gl}_N$ with basis $v_1, \dots, v_N$ such that $e_{i,j}v_k = \delta_{j,k}v_i$ for all i, j, k. Denote $V = (\mathbb{C}^N)^{\otimes n}$. For $I = (I_1, \dots, I_N) \in \mathcal{I}_n$, we define

$v_I \in V$ by the formula $v_I = v_{i_1} \otimes \cdots \otimes v_{i_n}$, where $i_j = i$ if $i_j \in I_i$. Let

$$V = \bigoplus_{|\boldsymbol{\lambda}|=n} V_{\boldsymbol{\lambda}}$$

be the weight decomposition. The vectors $(v_I)_{I \in \mathcal{I}_{\boldsymbol{\lambda}}}$ form a basis of $V_{\boldsymbol{\lambda}}$.

As always, we denote by $e^{(a)}_{i,j}$ the action of $e_{i,j}$ on the a-th tensor factor of V and denote $e_{i,j} = \sum_{a=1}^{n} e^{(a)}_{i,j}$.

Recall the $\widetilde{Y}(\mathfrak{gl}_N)$-module structure ϕ on $V \otimes \mathbb{C}[\boldsymbol{z};h]$ and the dynamical Hamiltonians X^q_i introduced in Section 4.1. The dynamical Hamiltonians preserve each of the weight subspaces $V_{\boldsymbol{\lambda}} \otimes \mathbb{C}[\boldsymbol{z};h]$.

Lemma 4.5 (Lemma 4.17 in [GRTV])**.** *For $i = 1, \dots, n$, we have*

$$\phi(X^q_i) = \sum_{a=1}^{n} z_a \, e^{(a)}_{i,i} + \frac{h}{2}(e^2_{i,i} - e_{i,i}) - h \sum_{j=1}^{N} \sum_{1 \leqslant a < b \leqslant n} e^{(a)}_{i,j} \, e^{(b)}_{j,i} - h \sum_{\substack{j=1 \\ j \neq i}}^{N} \frac{q_j}{q_i - q_j} \, G_{i,j} \,. \tag{4.8}$$

To obtain the lemma we replace h with $-h$ in Lemma 4.17 of [GRTV].

4.5. ***qKZ* difference connection.** Recall the $\widetilde{Y}(\mathfrak{gl}_N)$-action ϕ on $(\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}[\boldsymbol{z};h]$ introduced in Section 3.3. Let

$$R^{(i,j)}(u) = \frac{u - hP^{(i,j)}}{u - h}\,, \qquad i,j = 1, \dots, n\,, \quad i \neq j\,.$$

For $\kappa \in \mathbb{C}^\times$, define operators $K_1, \dots, K_n \in \mathrm{End}((\mathbb{C}^N)^{\otimes n}) \otimes \mathbb{C}[\boldsymbol{z};h]$,

$$\begin{aligned} K_i(\boldsymbol{z};\boldsymbol{q};h;\kappa) = {} & R^{(i+1,i)}(z_{i+1} - z_i) \, \dots \, R^{(n,i)}(z_n - z_i) \, \times \\ & \times \, q_1^{-e^{(i)}_{1,1}} \dots \, q_N^{-e^{(i)}_{N,N}} \, R^{(1,i)}(z_1 - z_i - \kappa) \, \dots \, R^{(i-1,i)}(z_{i-1} - z_i - \kappa)\,. \end{aligned}$$

Consider the difference operators $\widehat{K}_{\kappa,1}, \dots, \widehat{K}_{\kappa,n}$ acting on $(\mathbb{C}^N)^{\otimes n}$-valued functions of $\boldsymbol{z}, \boldsymbol{q}, h$,

$$\widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,i} F(\boldsymbol{z};\boldsymbol{q};h) = K_i(\boldsymbol{z};\boldsymbol{q};h;\kappa) \, F(z_1, \dots, z_{i-1}, z_i + \kappa, z_{i+1}, \dots, z_n; \boldsymbol{q}; h)\,. \tag{4.9}$$

Theorem 4.6 ([FR])**.** *The operators $\widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,1}, \dots, \widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,n}$ pairwise commute.*

Theorem 4.7 ([TV4])**.** *The operators $\widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,1}, \dots, \widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,n}$, $\phi(\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa,1}), \dots, \phi(\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa,N})$ pairwise commute.*

The commuting difference operators $\widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,1}, \dots, \widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,n}$ define the rational *qKZ* difference connection. We say that a $(\mathbb{C}^N)^{\otimes n}$-valued function $F(\boldsymbol{z};\boldsymbol{q};h)$ is a flat section of the difference connection if

$$\widehat{K}_{\boldsymbol{z},\boldsymbol{q},h,\kappa,i} F(\boldsymbol{z};\boldsymbol{q};h) = F(\boldsymbol{z};\boldsymbol{q};h), \qquad i = 1, \dots, n.$$

Theorem 4.7 says that the *qKZ* difference connection commutes with the trigonometric dynamical connection $\phi(\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa})$.

5. Yangian $\widetilde{Y}(\mathfrak{gl}_N)$ weight functions

5.1. **Weight functions W_I.** For $I \in \mathcal{I}_{\boldsymbol{\lambda}}$, we define the weight functions $W_I(\boldsymbol{t};\boldsymbol{z};h)$, c.f. [TV1, TV5].

Recall $\boldsymbol{\lambda} = (\lambda_1,\dots,\lambda_N)$. Denote $\lambda^{(i)} = \lambda_1 + \ldots + \lambda_i$ and $\lambda^{\{1\}} = \sum_{i=1}^{N-1}\lambda^{(i)} = \sum_{i=1}^{N-1}(N-i)\,\lambda_i$. Recall $I = (I_1,\dots,I_N)$. Set $\bigcup_{k=1}^{j} I_k = \{\, i^{(j)}_1 < \ldots < i^{(j)}_{\lambda^{(j)}} \}$. Consider the variables $t^{(j)}_a$, $j = 1,\dots,N$, $a = 1,\dots,\lambda^{(j)}$, where $t^{(N)}_a = z_a$, $a = 1,\dots,n$. Denote $t^{(j)} = (t^{(j)}_k)_{k\leqslant\lambda^{(j)}}$ and $\boldsymbol{t} = (t^{(1)},\dots,t^{(N-1)})$.

The weight functions are

$$W_I(\boldsymbol{t};\boldsymbol{z};h) \,=\, (-h)^{\lambda^{\{1\}}}\ \mathrm{Sym}_{t^{(1)}_1,\dots,t^{(1)}_{\lambda^{(1)}}} \,\ldots\, \mathrm{Sym}_{t^{(N-1)}_1,\dots,t^{(N-1)}_{\lambda^{(N-1)}}}\, U_I(\boldsymbol{t};\boldsymbol{z};h)\,, \tag{5.1}$$

$$U_I(\boldsymbol{t};\boldsymbol{z};h) \,=\, \prod_{j=1}^{N-1}\prod_{a=1}^{\lambda^{(j)}}\Biggl(\ \prod_{\substack{c=1\\ i^{(j+1)}_c < i^{(j)}_a}}^{\lambda^{(j+1)}} (t^{(j)}_a - t^{(j+1)}_c - h) \prod_{\substack{d=1\\ i^{(j+1)}_d > i^{(j)}_a}}^{\lambda^{(j+1)}} (t^{(j)}_a - t^{(j+1)}_d) \prod_{b=a+1}^{\lambda^{(j)}} \frac{t^{(j)}_a - t^{(j)}_b - h}{t^{(j)}_a - t^{(j)}_b}\Biggr)\,.$$

In these formulas for a function $f(t_1,\dots,t_k)$ of some variables we denote

$$\mathrm{Sym}_{t_1,\dots,t_k} f(t_1,\dots,t_k) = \sum_{\sigma\in S_k} f(t_{\sigma_1},\dots,t_{\sigma_k}).$$

Example. Let $N=2$, $n=2$, $\boldsymbol{\lambda} = (1,1)$, $I = (\{1\},\{2\})$, $J = (\{2\},\{1\})$. Then

$$W_I(\boldsymbol{t};\boldsymbol{z};h) = -h\,(t^{(1)}_1 - z_2), \qquad W_J(\boldsymbol{t};\boldsymbol{z};h) = -h\,(t^{(1)}_1 - z_1 - h).$$

5.2. **Weight functions $W_{\sigma,I}$.** For $\sigma\in S_n$ and $I \in \mathcal{I}_{\boldsymbol{\lambda}}$, we define

$$W_{\sigma,I}(\boldsymbol{t};\boldsymbol{z};h) = W_{\sigma^{-1}(I)}(\boldsymbol{t}; z_{\sigma(1)},\dots,z_{\sigma(n)};h),$$

where $\sigma^{-1}(I) = (\sigma^{-1}(I_1),\dots,\sigma^{-1}(I_N))$.

For a subset $A\subset\{1,\dots,n\}$, denote $\boldsymbol{z}_A = (z_a)_{a\in A}$. For $I\in\mathcal{I}_{\boldsymbol{\lambda}}$, denote $\boldsymbol{z}_I = (\boldsymbol{z}_{I_1},\dots,\boldsymbol{z}_{I_N})$. For $f(t^{(1)},\dots,t^{(N)}) \in \mathbb{C}[t^{(1)},\dots,t^{(N)}]^{S_{\lambda^{(1)}}\times\dots\times S_{\lambda^{(N)}}}$, we define $f(\boldsymbol{z}_I)$ by replacing $t^{(j)}$ with $\cup_{k=1}^{j}\boldsymbol{z}_{I_k}$. Denote

$$c_{\boldsymbol{\lambda}}(\boldsymbol{z}_I) \,=\, \prod_{a=1}^{N-1}\ \prod_{i,j\in\cup_{b=1}^{a} I_b} (z_i - z_j - h)\,,$$

$$R(\boldsymbol{z}_I) \,=\, \prod_{1\leqslant a<b\leqslant N}\ \prod_{i\in I_a}\prod_{j\in I_b}(z_i - z_j)\,, \qquad Q(\boldsymbol{z}_I) \,=\, \prod_{1\leqslant a<b\leqslant N}\ \prod_{i\in I_a}\prod_{j\in I_b}(z_i - z_j - h)\,.$$

5.3. **Stable envelope map.** Following [RTV], we define the weight function map

$$[W_{\mathrm{id}}] : V\otimes\mathbb{C}[\boldsymbol{z};h] \to H^*_T(\mathcal{X}_n), \quad v_I \mapsto [W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)],$$

where $W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)$ is the polynomial $W_{\mathrm{id},I}(\boldsymbol{t};\boldsymbol{z};h)$ in which variables $t^{(j)}_i$ are replaced with $\theta_{j,i}$ and $[W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)]$ is the cohomology class represented by $W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)$.

Denote

$$c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta}) = \prod_{a=1}^{N-1} \prod_{i=1}^{\lambda^{(a)}} \prod_{j=1}^{\lambda^{(a)}} (\theta_{a,i} - \theta_{a,j} - h) \in H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}}).$$

Observe that $c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta})$ is the equivariant Euler class of the bundle $\oplus_{a=1}^{N-1} \operatorname{Hom}(\boldsymbol{F}_a, \boldsymbol{F}_a)$ if we make $\mathbb{C}^\times$ act on it with weight $-h$. Note that $c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta})$ is not a zero-divisor in $H_T^*(\mathcal{X}_n)$, because none of its fixed point restrictions is zero.

Theorem 5.1 (Theorem 4.1 in [RTV]). *For any* $\boldsymbol{\lambda}$ *and any* $I \in \mathcal{I}_{\boldsymbol{\lambda}}$*, the cohomology class* $[W_{\mathrm{id},I}(\boldsymbol{\Theta}; \boldsymbol{z}; h)] \in H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *is divisible by* $c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta})$*, that is, there exists a unique element* $\mathrm{Stab}_{\mathrm{id},I} \in H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ *such that*

$$[W_{\mathrm{id},I}(\boldsymbol{\Theta}; \boldsymbol{z}; h)] \,=\, c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta}) \cdot \mathrm{Stab}_{\mathrm{id},I}\,. \tag{5.2}$$

By using this theorem we define the stable envelope map

$$\mathrm{Stab}_{\mathrm{id}} : V \otimes \mathbb{C}[\boldsymbol{z}; h] \to H_T^*(\mathcal{X}_n), \quad v_I \mapsto \mathrm{Stab}_{\mathrm{id},I}.$$

The stable envelope maps are main objects in [MO]. The stable envelope maps are defined in [MO] in terms of the torus T action on $\mathcal{X}_n$. Relation (5.2) gives a formula for the stable envelope map in terms of the Chern roots $\boldsymbol{\Theta}, \boldsymbol{z}, h$.

As we know, formula (3.3) defines the $\widetilde{Y}(\mathfrak{gl}_N)$-module structure ϕ on $V \otimes \mathbb{C}[\boldsymbol{z}; h]$, and formulas (3.4), (3.6), (3.7) define the $\widetilde{Y}(\mathfrak{gl}_N)$-module structure ρ on $H_T^*(\mathcal{X}_n)$.

Theorem 5.2 (Theorem 6.3 in [RTV]). *The stable envelope map* $\mathrm{Stab}_{\mathrm{id}} : V \otimes \mathbb{C}[\boldsymbol{z}; h] \to H_T^*(\mathcal{X}_n)$ *is a homomorphism of* $\widetilde{Y}(\mathfrak{gl}_N)$*-modules.*

5.4. **The inverse map.**

For $I \in \mathcal{I}_{\boldsymbol{\lambda}}$, introduce $\xi_I \in (\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}(\boldsymbol{z}; h)$ by the formula

$$\xi_I = \sum_{J \in \mathcal{I}_{\boldsymbol{\lambda}}} \frac{W_{\sigma_0,J}(\boldsymbol{z}_I; \boldsymbol{z}; h)}{Q(\boldsymbol{z}_I)\, c_{\boldsymbol{\lambda}}(\boldsymbol{z}_I)}\, v_J, \tag{5.3}$$

where $\sigma_0 \in S_n$ is the longest permutation.

Let $\mathbb{C}(\boldsymbol{z}; h)$ be the algebra of rational functions in $\boldsymbol{z}, h$. Consider the map

$$\nu = \oplus_{|\boldsymbol{\lambda}|=n} \nu_{\boldsymbol{\lambda}} \ : \ \oplus_{|\boldsymbol{\lambda}|=n} H_T^*(\mathcal{X}_{\boldsymbol{\lambda}}) \otimes \mathbb{C}(\boldsymbol{z}; h) \ \to \ (\mathbb{C}^N)^{\otimes n} \otimes \mathbb{C}(\boldsymbol{z}; h),$$

where $\nu_{\boldsymbol{\lambda}}$ is defined by the formula

$$[f(\boldsymbol{\Theta}; \boldsymbol{z}; h)] \mapsto \sum_{I \in \mathcal{I}_{\boldsymbol{\lambda}}} \frac{f(\boldsymbol{z}_I; \boldsymbol{z}; h)}{R(\boldsymbol{z}_I)} \xi_I\,,$$

see [GRTV, Formula (5.9)].

Lemma 5.3 (Lemma 6.7 in [RTV]). *The operator* ν *is inverse to* $\mathrm{Stab}_{\mathrm{id}}$.

5.5. **The difference connection on $H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$.**

By Theorem 4.7, the difference operators

$$\mathrm{Stab}_{\mathrm{id}} \circ \widehat{K}_{\boldsymbol{z}, \tilde{q}_1^{-1}, \dots, \tilde{q}_N^{-1}, -\kappa, 1} \circ \nu, \quad \dots, \quad \mathrm{Stab}_{\mathrm{id}} \circ \widehat{K}_{\boldsymbol{z}, \tilde{q}_1^{-1}, \dots, \tilde{q}_N^{-1}, -\kappa, n} \circ \nu$$

and the differential operators $\nabla_{\mathrm{quant}, \boldsymbol{\lambda}, \tilde{\boldsymbol{q}}, \kappa, 1}, \dots, \nabla_{\mathrm{quant}, \boldsymbol{\lambda}, \tilde{\boldsymbol{q}}, \kappa, N}$ pairwise commute. The difference operators form the rational *qKZ* difference connection on $H_T^*(T^*\mathcal{F}_{\boldsymbol{\lambda}})$. This difference connection is discussed in [MO] under the name of the shift operators.

6. Four more connections

In the remainder of this paper we fix $\boldsymbol{\lambda} \in \mathbb{Z}^N_{\geqslant 0}$, $|\boldsymbol{\lambda}| = n$.

6.1. **Trigonometric dynamical connection on $\pi_{V_{\boldsymbol{\lambda}}}$.** Consider $\mathbb{C}^n$ with coordinates $\boldsymbol{x} = (x_1, \dots, x_n)$ and $\mathbb{C}^N$ with coordinates $\boldsymbol{q} = (q_1, \dots, q_N)$. Consider the trivial bundle $\pi_{V_{\boldsymbol{\lambda}}} : V_{\boldsymbol{\lambda}} \times \mathbb{C}^{n+N} \to \mathbb{C}^{n+N}$.

Following [TV4] and [MTV1, Appendix B], introduce dynamical Hamiltonians

$$X^{V_{\boldsymbol{\lambda}}}_i(\boldsymbol{x}, \boldsymbol{q}) \,=\, \sum_{a=1}^{n} x_a\, e^{(a)}_{i,i} - \frac{e^2_{i,i}}{2} + \sum_{j=1}^{N} \sum_{1\leqslant a<b\leqslant n} e^{(a)}_{i,j}\, e^{(b)}_{j,i} + \sum_{\substack{j=1\\ j\neq i}}^{N} \frac{q_j}{q_i - q_j}\, G_{i,j}\,,$$

acting the $V_{\boldsymbol{\lambda}}$-valued functions of $\boldsymbol{x}, \boldsymbol{q}$. By formula (4.8), we have

$$\begin{aligned} \phi(X^q_i)(\boldsymbol{z}; \boldsymbol{q}; h) \,&=\, -hX^{V_{\boldsymbol{\lambda}}}_i(-z_1/h, \dots, -z_n/h; \boldsymbol{q}) - \frac{h}{2}\, e_{i,i} \,= \\ &=\, -hX^{V_{\boldsymbol{\lambda}}}_i(-z_1/h, \dots, -z_n/h; \boldsymbol{q}) - \frac{h}{2}\, \lambda_i\,. \end{aligned} \tag{6.1}$$

For $\kappa \in \mathbb{C}^{\times}$ the differential operators

$$\nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,i} \,=\, \kappa\, q_i \frac{\partial}{\partial q_i} - X^{V_{\boldsymbol{\lambda}}}_i(\boldsymbol{x}, \boldsymbol{q}), \qquad i = 1, \dots, N, \tag{6.2}$$

pairwise commute and define a flat connection $\nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa}$ on $\pi_{V_{\boldsymbol{\lambda}}}$, see [TV4].

6.2. **Rational *qKZ* difference connection on $\pi_{V_{\boldsymbol{\lambda}}}$.** Let

$$\tilde{R}^{(i,j)}(u) \,=\, \frac{u + P^{(i,j)}}{u+1}\,, \qquad i,j = 1, \dots, n\,, \quad i \neq j\,.$$

For $\kappa \in \mathbb{C}^{\times}$, define operators $K^{V_{\boldsymbol{\lambda}}}_1, \dots, \tilde{K}^{V_{\boldsymbol{\lambda}}}_n$ on $V_{\boldsymbol{\lambda}}$ by the formula

$$\begin{aligned} K^{V_{\boldsymbol{\lambda}}}_i(\boldsymbol{x}; \boldsymbol{q}; \kappa) \,=\, & \tilde{R}^{(i+1,i)}(x_{i+1} - x_i)\, \dots\, \tilde{R}^{(n,i)}(x_n - x_i)\, \times \\ & \times\, q_1^{-e^{(i)}_{1,1}} \dots\, q_N^{-e^{(i)}_{N,N}}\, \tilde{R}^{(1,i)}(x_1 - x_i - \kappa)\, \dots\, \tilde{R}^{(i-1,i)}(x_{i-1} - x_i - \kappa)\,. \end{aligned}$$

Consider the difference operators $K^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, K^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ acting on $(\mathbb{C}^N)^{\otimes n}$-valued functions of $\boldsymbol{x}, \boldsymbol{q}$,

$$\widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,i}\, F(\boldsymbol{x}, \boldsymbol{q}) \,=\, K^{V_{\boldsymbol{\lambda}}}_i(\boldsymbol{x}; \boldsymbol{q}; \kappa)\, F(x_1, \dots, x_{i-1}, x_i + \kappa, x_{i+1}, \dots, x_n; \boldsymbol{q})\,. \tag{6.3}$$

Theorem 6.1 ([FR])**.** *The operators* $\widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ *pairwise commute.*

Theorem 6.2 ([TV4])**.** *The operators* $\widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}\,,\ \nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,N}$ *pairwise commute.*

The commuting difference operators $\widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat{K}^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ define the rational *qKZ* difference connection on $\pi_{V_{\boldsymbol{\lambda}}}$. Theorem 6.2 says that the rational *qKZ* difference connection commutes with the trigonometric dynamical connection $\nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa}$.

6.3. **Relations between flat sections.**

Lemma 6.3. *Let $\kappa, h \in \mathbb{C}^\times$.*

(i) *Assume that $\tilde F(\boldsymbol{x};\boldsymbol{q}) \in V_{\boldsymbol{\lambda}}$ is a flat section of the connection $\nabla^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},-\kappa/h}$. Then*

$$F(\boldsymbol{z};\boldsymbol{q};h;\kappa) = \prod_{i=1}^N q_i^{-\frac{h\lambda_i}{2\kappa}} \prod_{1\leqslant i<j\leqslant N} (1-q_j/q_i)^{-\frac{h\,\min(\lambda_i,\lambda_j)}{\kappa}} \tilde F(-z_1/h,\dots,-z_n/h,\boldsymbol{q})$$

is a flat section of the connection $\phi(\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa})$.

(ii) *Assume that $\tilde F(\boldsymbol{x};\boldsymbol{q}) \in V_{\boldsymbol{\lambda}}$ is a flat section of the difference connection defined by the operators $\widehat K^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},-\kappa/h,1},\dots,\widehat K^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},-\kappa/h,n}$ in (6.3). Then $F(\boldsymbol{z};\boldsymbol{q};h)$ is a flat section of the difference connection defined by the operators $\widehat K_{\boldsymbol{z},\boldsymbol{q},\kappa,1},\dots,\widehat K_{\boldsymbol{z},\boldsymbol{q},\kappa,n}$ in (4.9).*

Proof. Part (i) follows from formulas (4.3) and (6.1). Part (ii) follows from formulas for the operators $\widehat K^{V_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},-\kappa/h,i}$ and $\widehat K_{\boldsymbol{z},\boldsymbol{q},\kappa,i}$. □

Lemma 6.4. *Assume $F(\boldsymbol{z};\tilde{\boldsymbol{q}},h) = \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} F_I(\boldsymbol{z};\tilde{\boldsymbol{q}},h)\,v_I \in V_{\boldsymbol{\lambda}}$ is a flat section of the connection $\phi(\nabla_{\boldsymbol{\lambda},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa})$ (resp. of the difference connection defined by the operators $\widehat K_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,1},\dots,\widehat K_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,n}$). Then*

$$\mathrm{Stab}_{\mathrm{id}}(F(\boldsymbol{z};\tilde{\boldsymbol{q}},h)) = \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} F_I(\boldsymbol{z};\tilde{\boldsymbol{q}};h)\,\mathrm{Stab}_{\mathrm{id},I} = \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} F_I(\boldsymbol{z};\tilde{\boldsymbol{q}};h)\,\frac{[W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)]}{c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta})} \tag{6.4}$$

is a flat section of the quantum connection $\nabla_{\mathrm{quant},\boldsymbol{\lambda},\tilde{\boldsymbol{q}},\kappa}$ (resp. of the difference connection defined by the operators $\mathrm{Stab}_{\mathrm{id}}\circ\widehat K_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,1}\circ\nu,\dots,\mathrm{Stab}_{\mathrm{id}}\circ\widehat K_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,n}\circ\nu$).

Proof. The first statement follows from Corollary 4.4. The second statement of obvious. □

6.4. **Trigonometric KZ connection on $W_{\boldsymbol{\lambda}}$.** Consider the Lie algebra $\mathfrak{gl}_n$. For $m\in\mathbb{Z}_{\geqslant 0}$, we denote by $W_m^{(n)}$ the irreducible $\mathfrak{gl}_n$-module with highest weight $(m,0,\dots,0)$ and highest weight vector w_m. For given $\boldsymbol{\lambda}=(\lambda_1,\dots,\lambda_N)$, $|\boldsymbol{\lambda}|=\lambda_1+\dots+\lambda_N=n$, we consider the $\mathfrak{gl}_n$-module $\otimes_{i=1}^N W_{\lambda_i}^{(n)}$. We denote by $W_{\boldsymbol{\lambda}}$ the weight subspace of $\otimes_{i=1}^N W_{\lambda_i}^{(n)}$ of $\mathfrak{gl}_n$-weight $(1,\dots,1)$. We consider the trivial bundle $\pi_{W_{\boldsymbol{\lambda}}}: W_{\boldsymbol{\lambda}}\times\mathbb{C}^{n+N}\to\mathbb{C}^{n+N}$.

Let $\Omega_0=\frac12\sum_{a=1}^n e_{a,a}\otimes e_{a,a}$, $\Omega_+=\Omega_0+\sum_{1\leqslant a<b\leqslant n} e_{a,b}\otimes e_{b,a}$, $\Omega_-=\Omega_0+\sum_{1\leqslant a<b\leqslant n} e_{b,a}\otimes e_{a,b}$. Introduce the trigonometric KZ operators $X_1^{W_{\boldsymbol{\lambda}}},\dots,X_N^{W_{\boldsymbol{\lambda}}}$ acting on $W_{\boldsymbol{\lambda}}$-valued functions of $\boldsymbol{x},\boldsymbol{q}$,

$$X_i^{W_{\boldsymbol{\lambda}}}(\boldsymbol{x};\boldsymbol{q}) = \sum_{a=1}^n\Big(x_a-\frac{e_{a,a}}{2}\Big)e_{a,a}^{(i)} + \sum_{\substack{j=1\\ j\neq i}}^N \frac{q_i\Omega_+^{(i,j)}+q_j\Omega_-^{(i,j)}}{q_i-q_j}\,.$$

For $\kappa\in\mathbb{C}^\times$, introduce the differential operators $\nabla^{W_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1},\dots,\nabla^{W_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,N}$,

$$\nabla^{W_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa,i} = \kappa q_i\frac{\partial}{\partial q_i} - X_i^{W_{\boldsymbol{\lambda}}}(\boldsymbol{x};\boldsymbol{q})\,. \tag{6.5}$$

The differential operators define on $\pi_{W_{\boldsymbol{\lambda}}}$ a flat connection $\nabla^{W_{\boldsymbol{\lambda}}}_{\boldsymbol{x},\boldsymbol{q},\kappa}$ called the trigonometric KZ connection.

6.5. **Rational dynamical difference connection on $\pi_{W_{\boldsymbol\lambda}}$.** For any $a, b = 1, \dots, n$, $a \neq b$, introduce a series $B_{a,b}(t)$ depending on a variable t,

$$B_{a,b}(t) = 1 + \sum_{s=1}^{\infty} (e_{b,a})^s (e_{a,b})^s \prod_{j=1}^{s} \frac{1}{j(t - e_{a,a} + e_{b,b} - j)} \,.$$

The series acts on any finite-dimensional $\mathfrak{gl}_n$-module W.

For $\kappa \in \mathbb{C}^\times$, introduce the operators $K_1^{W_{\boldsymbol\lambda}}, \dots, K_n^{W_{\boldsymbol\lambda}}$ acting on $W_{\boldsymbol\lambda}$-valued functions of $\boldsymbol{x}, \boldsymbol{q}$,

$$\begin{aligned} K_i^{W_{\boldsymbol\lambda}}(\boldsymbol{x}; \boldsymbol{q}; \kappa) \,=\, & \big(B_{i,n}(x_i - x_n) \,\dots\, B_{i,i+1}(x_i - x_{i+1})\big)^{-1} \times \\ & \times\, q_1^{-e^{(1)}_{i,i}} \dots\, q_N^{-e^{(N)}_{i,i}}\, B_{1,i}(x_1 - x_i - \kappa) \,\dots\, B_{i-1,i}(x_{i-1} - x_i - \kappa) \,. \end{aligned}$$

Introduce the difference operators $\widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$,

$$\widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,i} F(\boldsymbol{x}, \boldsymbol{q}) \,=\, K_i^{W_{\boldsymbol\lambda}}(\boldsymbol{x}, \boldsymbol{q}, \kappa)\, F(x_1, \dots, x_{i-1}, x_i + \kappa, x_{i+1}, \dots, x_n; \boldsymbol{q}). \tag{6.6}$$

Theorem 6.5 ([TV3])**.** *The operators* $\nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,N}$, $\widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ *pairwise commute.*

The commuting difference operators $\widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ define the rational dynamical difference connection on $\pi_{W_{\boldsymbol\lambda}}$. Theorem 6.5 says that the rational dynamical difference connection commutes with the trigonometric *KZ* connection $\nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa}$.

6.6. **Equivalence of connections on $\pi_{V_{\boldsymbol\lambda}}$ and $\pi_{W_{\boldsymbol\lambda}}$.** For $I = (I_1, \dots, I_N) \in \mathcal{I}_{\boldsymbol\lambda}$, define a vector $w_I \in W_{\boldsymbol\lambda}$ by the formula

$$w_I = w_{I_1} \otimes \dots \otimes w_{I_N}, \qquad w_{I_a} = \Big(\,{\prod_{i \in I_a}}' e_{i,1}\Big)\, w_{\lambda_a}, \tag{6.7}$$

where $\prod'$ means that we exclude from the product the factor $e_{1,1}$ if $1 \in I_a$. The map

$$\mu : W_{\boldsymbol\lambda} \to V_{\boldsymbol\lambda}, \qquad w_I \mapsto v_I, \tag{6.8}$$

is a vector isomorphism, see for example [TV4].

Theorem 6.6 (Theorem 5.8 in [TV4])**.** *The vector isomorphism μ identifies the action of operators* $\nabla^{V_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \nabla^{V_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$, $\widehat K^{V_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, \widehat K^{V_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$ *with the action of the operators* $\nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}$, $\dots$, $\nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$, $Q_1 \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1}, \dots, Q_n \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n}$, *respectively, where*

$$Q_i \,=\, \prod_{1 \leqslant j < i} \frac{x_j - x_i - 1 - \kappa}{x_j - x_i + 1 - \kappa} \prod_{i < j \leqslant n} \frac{x_i - x_j + 1}{x_i - x_j - 1}.$$

Corollary 6.7. *If a $W_{\boldsymbol\lambda}$-valued function $F^W(\boldsymbol{x}; \boldsymbol{q}) = \sum_{I \in \mathcal{I}_{\boldsymbol\lambda}} F^W_I(\boldsymbol{x}; \boldsymbol{q}) w_I$ satisfies the equations*

$$\nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1} F^W = 0, \quad \dots \quad \nabla^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,N} F^W = 0, \quad \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,1} F^W = F^W, \quad \dots \quad \widehat K^{W_{\boldsymbol\lambda}}_{\boldsymbol{x},\boldsymbol{q},\kappa,n} F^W = F^W,$$

then the $V_{\boldsymbol\lambda}$-valued function

$$F^V(\boldsymbol x;\boldsymbol q) = \prod_{1\leqslant i<j\leqslant n}\frac{\Gamma((x_i-x_j-1)/\kappa)}{\Gamma((x_i-x_j+1)/\kappa)}\sum_{I\in\mathcal I_{\boldsymbol\lambda}} F^W_I(\boldsymbol x;\boldsymbol q)\,v_I$$

satisfies the equations

$$\nabla^{V_{\boldsymbol\lambda}}_{\boldsymbol x,\boldsymbol q,\kappa,1}F^V=0,\quad \dots\quad \nabla^{V_{\boldsymbol\lambda}}_{\boldsymbol x,\boldsymbol q,\kappa,N}F^V=0,\quad \widehat K^{V_{\boldsymbol\lambda}}_{\boldsymbol x,\boldsymbol q,\kappa,1}F^V=F^V,\quad \dots\quad \widehat K^{V_{\boldsymbol\lambda}}_{\boldsymbol x,\boldsymbol q,\kappa,n}F^V=F^V.$$

7. Integral representations for flat sections

7.1. **Master function.** Consider $\mathbb C^{\frac{n(n-1)}{2}}$ with coordinates $\boldsymbol s = (s_j^{(a)})$, $a=1,\dots,n-1$, $j=1,\dots,n-a$. Consider $\mathbb C^n$ with coordinates $\boldsymbol x=(x_1,\dots,x_n)$ and $\mathbb C^N$ with coordinates $\boldsymbol q=(q_1,\dots,q_N)$. The master function is

$$\Phi_{\boldsymbol\lambda}(\boldsymbol s;\boldsymbol x;\boldsymbol q) = \prod_{1\leqslant i<j\leqslant N}(q_i-q_j)^{\lambda_i\lambda_j}\prod_{i=1}^{N} q_i^{\lambda_i(x_1-1+\lambda_i/2)}\prod_{j=1}^{n-1}\prod_{i=1}^{N}(s_j^{(1)}-q_i)^{-\lambda_i}\times$$
$$\times\prod_{a=1}^{n-1}\prod_{i=1}^{n-a}(s_i^{(a)})^{x_{a+1}-x_a+1}\prod_{a=1}^{n-2}\Big(\prod_{1\leqslant i<j\leqslant n-a}(s_i^{(a)}-s_j^{(a)})^2\prod_{i=1}^{n-a}\prod_{j=1}^{n-a-1}(s_i^{(a)}-s_j^{(a+1)})^{-1}\Big),$$

see [MV, Formula (16)].

7.2. $\mathfrak{gl}_n$ **weight functions.** For $I\in\mathcal I_{\boldsymbol\lambda}$ we define the weight function $\omega_I(\boldsymbol s;\boldsymbol q)$, see [MV, RSV, SV]. Introduce new variables $\hat{\boldsymbol s}=(s_{a,i,j})$, where $a\in\{1,\dots,N\}$, $i\in I_a$, $j\in\{1,\dots,i-1\}$. For $k=1,\dots,n-1$, denote by $\hat{\boldsymbol s}_k$ the set of all variables $s_{a,i,j}$ with $j=k$. Then $|\hat{\boldsymbol s}_k|=n-k$.

Let B be the set of sequences $\boldsymbol\beta=(\beta_1,\dots,\beta_{n-1})$ of bijections $\beta_k:\hat{\boldsymbol s}_k\to\{s_1^{(k)},\dots,s_{n-k}^{(k)}\}$, $k=1,\dots,n-1$.

If $f(\hat{\boldsymbol s})$ is a function of $\hat{\boldsymbol s}$ and $\boldsymbol\beta\in B$, then we obtain the function $(\boldsymbol\beta f)(\boldsymbol s)$ of $\boldsymbol s$ by replacing variables in $f(\hat{\boldsymbol s})$ according to the bijection $\boldsymbol\beta$.

Introduce the function

$$f_I(\hat{\boldsymbol s};\boldsymbol q)=\prod_{a=1}^{N}\prod_{i\in I_a}\frac{1}{(s_{a,i,1}-q_a)(s_{a,i,2}-s_{a,i,1})\dots(s_{a,i,i-1}-s_{a,i,i-2})},$$

where we set $\frac{1}{(s_{a,i,1}-q_a)(s_{a,i,2}-s_{a,i,1})\dots(s_{a,i,i-1}-s_{a,i,i-2})}=1$ if $i=1$. We define

$$\omega_I(\boldsymbol s;\boldsymbol q)=\sum_{\boldsymbol\beta\in B}(\boldsymbol\beta f_I)(\boldsymbol s;\boldsymbol q).$$

Example. Let $N=2$, $n=3$, $\boldsymbol\lambda=(1,2)$, $I=(I_1,I_2)$, $I_1=\{2\}$, $I_2=\{1,3\}$. Then

$$\omega_I(\boldsymbol s;\boldsymbol q)=\frac{1}{(s_1^{(1)}-q_1)(s_2^{(1)}-q_2)(s_1^{(2)}-s_2^{(1)})}+\frac{1}{(s_2^{(1)}-q_1)(s_1^{(1)}-q_2)(s_1^{(2)}-s_1^{(1)})}.$$

7.3. **Hypergeometric integrals.** In the space $\mathbb{C}^{\frac{n(n-1)}{2}} \times \mathbb{C}^n \times \mathbb{C}^N$ with coordinates $\boldsymbol{s}, \boldsymbol{x}, \boldsymbol{q}$ we consider the arrangement $\mathcal{C}$ of hyperplanes defined by equations

$$\begin{aligned}
& s_i^{(a)} = 0, \qquad a = 1, \dots, n-1, \quad i = 1, \dots, n-a; \\
& s_j^{(1)} - q_i = 0, \qquad j = 1, \dots, n-1, \quad i = 1, \dots, N; \\
& s_i^{(a)} - s_j^{(a)} = 0, \qquad a = 1, \dots, n-1, \quad 1 \leqslant i < j \leqslant n-a; \\
& s_i^{(a)} - s_j^{(a+1)} = 0, \qquad a = 1, \dots, n-2, \quad i = 1, \dots, n-a, \quad j = 1, \dots, n-a-1. \\
& q_i - q_j = 0, \qquad 1 \leqslant i < j \leqslant N; \\
& q_i = 0, \qquad i = 1, \dots, N.
\end{aligned}$$

Denote by U the complement to the union of hyperplanes of the arrangement $\mathcal{C}$.

In the space $\mathbb{C}^n \times \mathbb{C}^N$ with coordinates $\boldsymbol{x}, \boldsymbol{q}$ consider the arrangement of hyperplanes defined by equations

$$\begin{aligned}
& q_i - q_j = 0, \qquad 1 \leqslant i < j \leqslant N; \\
& q_i = 0, \qquad i = 1, \dots, N.
\end{aligned}$$

Denote by Δ the complement to the union of hyperplanes of this arrangement.

Consider the projection $\pi : \mathbb{C}^{\frac{n(n-1)}{2}} \times \mathbb{C}^n \times \mathbb{C}^N \to \mathbb{C}^n \times \mathbb{C}^N$. For every $(\boldsymbol{x}; \boldsymbol{q})$ the arrangement $\mathcal{C}$ induces an arrangement in the fiber of π over $(\boldsymbol{x}; \boldsymbol{q})$. Denote by $U(\boldsymbol{x}; \boldsymbol{q})$ the complement to that arrangement in the fiber.

Consider the master function $\Phi_{\boldsymbol{\lambda}}(\boldsymbol{s}; \boldsymbol{x}; \boldsymbol{q})$ as a multivalued function on U. Let $\kappa \in \mathbb{C}^\times$. The function $\Phi_{\boldsymbol{\lambda}}(\boldsymbol{s}; \boldsymbol{x}; \boldsymbol{q})^{1/\kappa}$ defines a rank one local system $\mathcal{L}_\kappa$ on U whose horizontal sections over open subsets of U are univalued branches of $\Phi_{\boldsymbol{\lambda}}(\boldsymbol{s}; \boldsymbol{x}; \boldsymbol{q})^{1/\kappa}$ multiplied by complex numbers. The vector bundle

$$\sqcup_{(\boldsymbol{x}, \boldsymbol{q}) \in \Delta} \; H_{\frac{n(n-1)}{2}}(U(\boldsymbol{x}; \boldsymbol{q}), \mathcal{L}_\kappa |_{U(\boldsymbol{x}; \boldsymbol{q})}) \to \Delta \tag{7.1}$$

has the canonical flat Gauss-Manin connection. Let

$$\psi(\boldsymbol{x}, \boldsymbol{q}, \kappa) \in H_{\frac{n(n-1)}{2}}(U(\boldsymbol{x}, \boldsymbol{q}), \mathcal{L}_\kappa |_{U(\boldsymbol{x}, \boldsymbol{q})})$$

be a flat section of the Gauss-Manin connection. In what follows we will consider the multidimensional hypergeometric integrals

$$\mathcal{J}_{\psi, I}(\boldsymbol{x}; \boldsymbol{q}; \kappa) = \int_{\psi(\boldsymbol{x}, \boldsymbol{q}, \kappa)} \Phi_{\boldsymbol{\lambda}}(\boldsymbol{s}; \boldsymbol{x}; \boldsymbol{q})^{1/\kappa} \omega_I(\boldsymbol{s}; \boldsymbol{q}) \wedge_{a=1}^{n-1} \wedge_{i=1}^{n-a} ds_i^{(a)}$$

for $I \in \mathcal{I}_{\boldsymbol{\lambda}}$.

Theorem 7.1 ([MV]). *For every flat section $\psi(\boldsymbol{x}, \boldsymbol{q}, \kappa)$ of the Gauss-Manin connection on the bundle in* (7.1), *the function*

$$F_\psi(\boldsymbol{x}; \boldsymbol{q}; \kappa) = \sum_{I \in \mathcal{I}_{\boldsymbol{\lambda}}} \mathcal{J}_{\psi, I}(\boldsymbol{x}; \boldsymbol{q}; \kappa) \, w_I \tag{7.2}$$

is a flat section of the trigonometric KZ connection (6.5) *on $\pi_{W_{\boldsymbol{\lambda}}}$ and of the rational dynamical difference connection* (6.6) *on $\pi_{W_{\boldsymbol{\lambda}}}$.*

Corollary 7.2. *The $V_{\boldsymbol{\lambda}}$-valued function*

$$F^V_\psi(\boldsymbol{x};\boldsymbol{q};\kappa) = \prod_{1\leqslant i<j\leqslant n} \frac{\Gamma((x_i-x_j-1)/\kappa)}{\Gamma((x_i-x_j+1)/\kappa)} \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} \mathcal{J}_{\psi,I}(\boldsymbol{x};\boldsymbol{q};\kappa)\, v_I$$

is a flat section of the trigonometric dynamical connection (6.2) *on $\pi_{V_{\boldsymbol{\lambda}}}$ and of the rational qKZ difference connection* (6.3) *on $\pi_{V_{\boldsymbol{\lambda}}}$.*

Proof. The corollary follows from Corollary 6.7. □

Corollary 7.3. *The $V_{\boldsymbol{\lambda}}$-valued function*

$$\begin{aligned} F_\psi(\boldsymbol{z};\boldsymbol{q};h;\kappa) &= \prod_{i=1}^N q_i^{-\frac{h\lambda_i}{2\kappa}} \prod_{1\leqslant i<j\leqslant N} (1-q_j/q_i)^{-\frac{h\,\min(\lambda_i,\lambda_j)}{\kappa}} \times \\ &\times \prod_{1\leqslant i<j\leqslant n} \frac{\Gamma((z_i-z_j+h)/\kappa)}{\Gamma((z_i-z_j-h)/\kappa)} \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} \mathcal{J}_{\psi,I}(-z_1/h,\dots,-z_n/h;\boldsymbol{q};-\kappa/h)\, v_I \end{aligned}$$

is a flat section of the trigonometric dynamical connection $\phi(\nabla_{\boldsymbol{\lambda},\boldsymbol{q},\kappa})$ on $\pi_{V_{\boldsymbol{\lambda}}}$ and of the difference connection on $\pi_{V_{\boldsymbol{\lambda}}}$ defined by the operators $\widehat{K}_{\boldsymbol{z},\boldsymbol{q},\kappa,1},\dots,\widehat{K}_{\boldsymbol{z},\boldsymbol{q},\kappa,n}$ in (4.9)*.*

Proof. The corollary follows from Lemma 6.3. □

Corollary 7.4. *For any ψ as in Section 7.3,*

(7.3)
$$\begin{aligned} &\mathrm{Stab}_{\mathrm{id}}(F_\psi(\boldsymbol{z};\tilde q_1^{-1},\dots,\tilde q_N^{-1};h;-\kappa)) = \\ &= \prod_{i=1}^N \tilde q_i^{-\frac{h\lambda_i}{2\kappa}} \prod_{1\leqslant i<j\leqslant N} (1-\tilde q_i/\tilde q_j)^{\frac{h\,\min(\lambda_i,\lambda_j)}{\kappa}} \prod_{1\leqslant i<j\leqslant n} \frac{\Gamma((z_j-z_i-h)/\kappa)}{\Gamma((z_j-z_i+h)/\kappa)} \\ &\quad\times \sum_{I\in\mathcal{I}_{\boldsymbol{\lambda}}} \mathcal{J}_{\psi,I}(-z_1/h,\dots,-z_n/h;\tilde q_1^{-1},\dots,\tilde q_N^{-1};\kappa/h)\, \frac{[W_{\mathrm{id},I}(\boldsymbol{\Theta};\boldsymbol{z};h)]}{c_{\boldsymbol{\lambda}}(\boldsymbol{\Theta})} \end{aligned}$$

is a flat section of the quantum connection $\nabla_{\mathrm{quant},\boldsymbol{\lambda},\tilde{\boldsymbol{q}},\kappa}$ on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ and of the difference connection on $H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}})$ defined by the operators $\mathrm{Stab}_{\mathrm{id}}\circ \widehat{K}_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,1}\circ\nu,\dots,$ $\mathrm{Stab}_{\mathrm{id}}\circ \widehat{K}_{\boldsymbol{z},\tilde q_1^{-1},\dots,\tilde q_N^{-1},-\kappa,n}\circ\nu$.

Proof. The corollary follows from Lemma 6.4. □

7.4. **Example.** Let $n=N=2$, $\boldsymbol{\lambda}=(1,1)$. Then $T^*\mathcal{F}_{\boldsymbol{\lambda}}$ is the cotangent bundle of projective line and

$$H^*_T(T^*\mathcal{F}_{\boldsymbol{\lambda}}) = \mathbb{C}[\gamma_{1,1},\gamma_{2,1}]\otimes\mathbb{C}[z_1,z_2]\otimes\mathbb{C}[h]/\langle (u-\gamma_{1,1})(u-\gamma_{2,1}) = (u-z_1)(u-z_2)\rangle.$$

We have

$$\begin{aligned} \mathrm{Stab}_{\mathrm{id}}(F_\psi(\boldsymbol{z};\tilde q_1^{-1},\tilde q_2^{-1};h;-\kappa)) &= \tilde q_1^{(z_1+h)/\kappa}\tilde q_2^{z_1/\kappa}(\tilde q_1^{-1}-\tilde q_2^{-1})^{2h/\kappa}\frac{\Gamma((z_2-z_1-h)/\kappa)}{\Gamma((z_2-z_1+h)/\kappa)} \times \\ &\times \int_\psi s^{(z_1-z_2+h)/\kappa}(s-\tilde q_1^{-1})^{-h/\kappa}(s-\tilde q_2^{-1})^{-h/\kappa}\Big[\frac{\gamma_{1,1}-z_1-h}{s-\tilde q_1^{-1}} + \frac{\gamma_{1,1}-z_2}{s-\tilde q_2^{-1}}\Big]\, ds. \end{aligned}$$

The master function is

$$\Phi(s;\boldsymbol{z};\boldsymbol{q};h)=s^{(z_1-z_2+h)/\kappa}(s-\tilde q_1^{-1})^{-h/\kappa}(s-\tilde q_2^{-1})^{-h/\kappa}.$$

The critical point equation is

$$\frac{z_1-z_2+h}{s}-\frac{h}{s-\tilde q_1^{-1}}-\frac{h}{s-\tilde q_2^{-1}}=0$$

or

$$(z_2-z_1+h)s^2-(z_2-z_1)(\tilde q_1^{-1}+\tilde q_2^{-1})s+\tilde q_1^{-1}\tilde q_2^{-1}(z_2-z_1-h)=0.$$

The discriminant of this equation is

$$\begin{aligned}\Delta_\Phi(\boldsymbol{z};\tilde{\boldsymbol{q}};h)&=(z_1-z_2)^2(\tilde q_1^{-1}-\tilde q_2^{-1})^2+4\tilde q_1^{-1}\tilde q_2^{-1}h^2=\\&=\frac{(z_1-z_2)^2(\tilde q_1-\tilde q_2)^2+4\tilde q_1\tilde q_2h^2}{\tilde q_1^2\tilde q_2^2}\,.\end{aligned}\tag{7.4}$$

On the other hand, the algebra of quantum multiplication $\mathcal{H}^q_{\boldsymbol\lambda}$ is the quotient of $\mathbb{C}[\gamma_{1,1},\gamma_{2,1},\boldsymbol{z};h]$ by the relations

$$\gamma_{1,1}+\gamma_{2,1}\,=\,z_1+z_2\,,\qquad \gamma_{1,1}\gamma_{2,1}+\frac{\tilde q_1}{\tilde q_1-\tilde q_2}\,h\,(\gamma_{1,1}-\gamma_{2,1}-h)\,=\,z_1z_2\,,$$

see the example in [GRTV, Section 6] where again we replaced h with $-h$ and q_1,q_2 with $\tilde q_1^{-1},\tilde q_2^{-1}$, respectively. Set $\gamma_{1,1}=t,\gamma_{2,1}=z_1+z_2-t$. Then the relation is

$$(\tilde q_1-\tilde q_2)t^2-(2\tilde q_1h+(\tilde q_1-\tilde q_2)(z_1+z_2))t+(\tilde q_1-\tilde q_2)z_1z_2+\tilde q_1h(z_1+z_2+h)=0.$$

The discriminant of this equation is

$$\Delta_{\mathcal{H}^q_{\boldsymbol\lambda}}(\boldsymbol{z};\tilde{\boldsymbol{q}};h)=(z_1-z_2)^2(\tilde q_1-\tilde q_2)^2+4\tilde q_1\tilde q_2h^2.\tag{7.5}$$

By comparing (7.4) and (7.5), we conclude that the algebra of functions on the critical set of the master function is isomorphic to the algebra of quantum multiplication.

References

[BMO] A. Braverman, D. Maulik, A. Okounkov, *Quantum cohomology of the Springer resolution*, Preprint (2010), 1–35, `arXiv:1001.0056`

[FR] I. Frenkel, N. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), no. 1, 1–60

[G1] A. Givental, *Equivariant Gromov-Witten invariants*, Internat. Math. Res. Notices (1996), no. 13, 613-663

[G2] A. Givental, *Stationary phase integrals, quantum Toda lattices, flag manifolds and the mirror conjecture*, Topics in singularity theory, AMS Transl. Ser. 2 **180** (1997), 103–115

[G3] A. Givental, *Gromov-Witten invariants and quantization of quadratic Hamiltonians*, Mosc. Math. J. **1** (2001), no. 4, 551–568

[MO] D. Maulik, A. Okounkov, *Quantum Groups and Quantum Cohomology*, Preprint (2012), 1–276, `arXiv:1211.1287`

[GRTV] V. Gorbounov, R. Rimanyi, V. Tarasov, A. Varchenko, *Cohomology of the cotangent bundle of a flag variety as a Yangian Bethe algebra*, `arXiv:1204.5138`, 1–44

[MTV1] E. Mukhin, V. Tarasov, A. Varchenko, *Bethe eigenvectors of higher transfer matrices*, J. Stat. Mech. (2006), no. 8, P08002, 1–44

[MTV2] E. Mukhin, V. Tarasov, A. Varchenko, *Bethe algebra of the $\mathfrak{gl}_{N+1}$ Gaudin model and algebra of functions on the critical set of the master function*, New trends in quantum integrable systems, 307–324, World Sci. Publ., Hackensack, NJ, 2011

[MTV3] E. Mukhin, V. Tarasov, A. Varchenko, *Three sides of the geometric Langlands correspondence for $\mathfrak{gl}_N$ Gaudin model and Bethe vector averaging maps*, Arrangements of hyperplanes, Sapporo 2009, 475–511, Adv. Stud. Pure Math., 62, Math. Soc. Japan, Tokyo, 2012

[MV] Y. Markov and A. Varchenko, *Hypergeometric solutions of trigonometric KZ equations satisfy dynamical difference equations*, Adv. Math. **166** (2002), no. 1, 100–147

[N1] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac- Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416

[N2] H. Nakajima, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3, 515–560

[RSV] R. Rimanyi, L. Stevens, A. Varchenko, *Combinatorics of rational functions and Poincare-Birchoff-Witt expansions of the canonical $U(\mathfrak{n}_-)$-valued differential form*, Ann. Comb. **9** (2005), no. 1, 57–74

[RTV] R. Rimanyi, V. Tarasov, A. Varchenko, *Partial flag varieties, stable envelopes and weight functions*, Preprint (2012), 1–24, `arXiv:1212.6240`

[SV] V. Schechtman and A. Varchenko, *Arrangements of hyperplanes and Lie algebra homology*, Invent. Math. **106** (1991), 139–194

[TV1] V. Tarasov and A. Varchenko, *Jackson integral representations for solutions of the quantized Knizhnik-Zamolodchikov equation*, (Russian) Algebra i Analiz 6 (1994), no. 2, 90–137; translation in St. Petersburg Math. J. 6 (1995), no. 2, 275–313

[TV2] V. Tarasov and A. Varchenko, *Geometry of q-hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), no. 3, 501–588

[TV3] V. Tarasov and A. Varchenko, *Difference equations compatible with trigonometric KZ differential equations*, Internat. Math. Res. Notices 2000, no. 15, 801–829

[TV4] V. Tarasov and A. Varchenko, *Duality for Knizhnik-Zamolodchikov and dynamical equations*, Preprint (2001), 1–12, `arXiv:math/0112005`

[TV5] V. Tarasov and A. Varchenko, *Combinatorial formulae for nested Bethe vectors*, Preprint (2007), 1–31, `arXiv:math/0702277`

[TL] V. Toledano Laredo, *The trigonometric Casimir connection of a simple Lie algebra*, J. Algebra **329** (2011), 286–327